arXiv:1301.3238v1 [nlin.AO] 15 Jan 2013

Cycle frequency in standard Rock-Paper-Scissors games: Evidence from experimental economics

Bin Xu[1,2,3] and Zhijian Wang[2,3*]
[1] Public Administration College, Zhejiang Gongshang University, Hangzhou, 310018, China
[2] Experimental Social Science Laboratory, Zhejiang University, Hangzhou, 310058, China and
[3] State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,
Chinese Academy of Sciences, Beijing, 100190, China
(Dated: January 16, 2013)

In the iconic game, standard Rock-Paper-Scissors (RPS) game, cycle in evolutionary trajectory is predicted by evolutionary game theory but not by classical game theory. We conduct lab experiments of the standard RPS game to index the cycle. Our experiments are of traditional setting with discrete time and random matched. In the 12 samples of 300 rounds 'stochastic' social evolution trajectories, (1) We confirm that the cycles not only exist but also persist; (2) We firstly test out the frequencies of cycles in this game; The observed frequency is about 0.028 ± 0.007 period per experimental round. The application of the observed frequency on a dynamics model is illustrated. Using frequency of cycles to index the distance from equilibrium, we consider both of the time symmetry and the sampling size effect, both of which have the root in non-equilibrium statistical physics.

PACS numbers: 89.65.-s 87.23.Cc 02.50.Le 01.50.My

INTRODUCTION

In evolutionary trajectories, formulating a physically meaningful measure of the distance from equilibrium is itself an area of active research [1]. An equilibrium system satisfies detailed balance condition, in which time reversal symmetry holds and the distance from equilibrium is zero. If a system is in non-equilibrium stationary state, the long-run balances of transits within states are broken and the distance from equilibrium should deviate from zero. Observations, e.g. entropy production [2] and velocity [3], have been developed to index the distance from equilibrium. This picture comes from non-equilibrium statistical physics. In this paper, this picture is borrowed to investigate the trajectories from lab experiments in game theory.

Game theory studies strategies interactions and social evolutionary trajectories in the systems from bacteria to human being [4–7]. In game theory, there exist two theories — classical game theory (CGT) [8] and evolutionary game theory (EGT) [9]. From the two theories, the predictions are identical on Nash equilibrium, but different on the evolutionary trajectories.

We use the standard Rock-Paper-Scissors (RPS) game as an example. This game is iconic in both CGT and EGT [8–10]. In the evolutionary trajectories surround Nash equilibrium, in long run, EGT predicts the existence of cycles, the system is in non-equilibrium and the transits within the states are not in detailed balance. Alternatively, CGT predict full random — the system is in equilibrium, in long run, each loop should be balanced by its reverse loop. Because, the system is in equilibrium state, in which corresponding to every individual process there is a reverse process, and then the average rate of every process is equal to the average rate of its reverse process [11]; So CGT predicts that, in long run, *excess loops* (hereafter, called as cycles) cannot be observed in the trajectories. So, the crucial bifurcation of the two theories is whether the cycles are existent. Using cycles as an object, we will show that the distance from equilibrium can be measured, and then can be employed to evaluate evolutionary dynamics model. We carry out these tasks in lab experiments.

Lab experiments study human subjects' strategy interactions and have been major tool to test game theory [12]. In past decades, lab game experiments have been employed to test EGT [13], in which to report the cycles is a task [14–16]. The existence of cycles in the evolutionary trajectory has been supported only with figures qualitatively [14, 15]. Only quite recently, in RPS-like game, but in a special setting (continuous time) games, the existence cycles are firstly observed [17]. Regrettably, in traditional setting (discrete time and random matching [13–15, 18–21]) game experiment environments, in the evolutionary trajectories, cycles has never been indexed quantitatively.

Discrete time games, rooting in the natural period (e.g., the year for animal [5]), are concerned by experimenters [10, 13–15, 18–21] and theorists [22, 23]. Our experiments are standard RPS games of discrete time. In these experiments, we harvest 12 experimental trajectories and each includes 300 rounds. Like in existed experiments [14, 15, 18, 20, 21], the evolutionary trajectories are high stochastic. Borrowing the picture from non-equilibrium statistical physics to measure the distance from equilibrium (or detailed balance) in trajectories [24], we index the cycles (excess loops surround

---

*Corresponding author, email:wangzj@zju.edu.cn

	R	P	S
R	1	0	2
P	2	1	0
S	0	2	1

FIG. 1: Payoff matrix of the standard RPS Game.

Nash equilibrium) in the stochastic evolutionary trajectories quantitatively.

In our discrete time standard RPS game experiments, we confirm the cycles exist, persist and do not dissipate; Then, we test out the observed frequency of the cycles is about 0.028 (±0.007) period per experimental round. Then, we demonstrate how to evaluate an EGT model with the observed frequency.

This paper is organized as following. In next session, we introduce the traditional setting standard RPS experiment. In section 3, the measurement methods are introduced; In section 4, we report the performance of the cycles and the support materials; Section 5 is to demonstrate also how to evaluate models with frequency. Discussion and summary is in the last.

EXPERIMENT

In RPS game, rock (R) beats scissors (S) which beat paper (P) which beats rock. Standard RPS means, taking tie as reference, the benefit for winner is equal to loss for the opponent. This game is simplest and iconic for both CGT and EGT [10, 17, 25]. The payoff matrix used in the experiment is presented in Fig. 1. One will get 2 experiment points if win, 1 if tie and 0 if loss.

The laboratory experiments consist of 12 independent groups. Each group consists of 6 players; so our experiment is a finite population (the population size is 6) game experiment. Each group plays the RPS game 300 rounds (the aim of this 300-round setting should be discussed following). In each experimental round, every player plays the game against a opponent who is randomly selected by computer program within a group. Such experimental setting is a traditional in experimental economics [13–15, 18–21]. All subjects make their own decisions simultaneously and anonymously. Only after all subjects submitted their choices, each of them would get the feedback including her/his strategy, her/his opponent's strategy and her/his own point through their private screen. No other information was provided. And these are common knowledge.

The 12 experimental sessions were conducted during Dec. 2010 in experimental social science laboratory, Zhejiang University. All 72 subjects were recruited broadly from the student populations at Zhejiang University. They were sitting in an isolated seat with a computer after arrived. Both written and oral instructions were provided for every subject before experiment implementing. During the experiment, subjects gained experiment points each round according to the payoff matrix. The experimental sessions were lasted about 1.5 to 2 hours. The subjects got the earnings in cash privately after the experiments according to the accumulated points over 300 experimental rounds. The exchange rule is 1 experiment point equals 0.15 Yuan RMB. In addition, each subject get 5 Yuan RMB as show-up fee. The average earning is 50 Yuan RMB.

NOTIONS AND MEASUREMENT

Notions

Social state —— Social state has been used to represent social performance in game widely [10, 26]. In evolutionary game theory view, the social state is described by the densities of strategies in a society. In RPS game, the strategy set is (S, R, P), so the social state is denoted as (x, y, z), herein, x, y, z are the densities of strategy Scissors, Rock and Paper in a society respectively. Since $z = 1 - x - y$, the state can be represented in simply as (x, y). If there are six persons in a group (society) just as our experimental setting, the total number of state is 28.

State space—— All potential states compose the state space. In the case of our six person RPS game, the state space is an equilateral triangle including 28 states in a two dimensional plane. We illustrate the state space and 28 social states in Fig. 2. For example, state $(0, 0)$ in the bottom left represents there are 0 scissors, 0 rock and 6 paper in the group, and the state $(1/3, 1/3)$ in the center represents 2 scissors, 2 rock and 2 paper in the group.

Evolutionary trajectory —— Evolutionary trajectory is the evolutionary trace of social state transit in state space [10, 14, 17, 26]. In our experiments, each six person group repeated the game 300 times. For each round, the social state should be in and only in one of the discrete 28 states. From round t to $t+1$, the social state may change and we call this change as transit. The social state should transit 299 times during an experiment and form an evolutionary trajectory.

Cycles in a trajectory —— In the view of evolutionary game theory, the evolutionary trajectory in RPS game here should form counter-clockwise cycles [9, 10, 25]. However, the evolutionary trajectory is full of noises and we can hardly test out robust pattern, especially in short run. Strictly speaking, in our experimental trajectories, the cycles are the observed loops of transits



FIG. 2: State space and Poincare section. There are total 28 observable states in six person RPS game. The state space is an equilateral triangle including 28 states in two dimensional plane, in which, state $(0,0)$ in the bottom left represents there are 0 scissors, 0 rock and 6 paper in the group, and the state $(1/3,1/3)$ in the center represents 2 scissors, 2 rock and 2 paper in the group. The segment S is the Poincare section for cycles counting. In our measurement, the segment is shifted (to left-down in 10^{-3}) slightly and it does not pass through any of the 28-state.

surround N_e in one direction *excess* over the opposite direction [1, 2, 24].

Distance from equilibrium —— One point need to be clarified. The distance from equilibrium in this paper is to index the asymmetry, or unbalance, of the transitions within states in trajectories. As mentioned, our distance from equilibrium is indexed by the cycles (the excess loops) in the trajectory. On the contrast, in game experiment literatures, e.g., [10, 17] which are most close to this investigate, the distance from equilibrium, called as the distance from the center [10, 17], is to index the distance from an observed state to a Nash equilibrium point.

Measurement

Poincare section and cycles —— Following Ref [24] using excess transitions to represent the deterministic process in stochastic processes, we further use cycle counting method to represent the cycles and take it as the index of distance from equilibrium. Poincare section which introduced into experimental economics by Cason et al [17] is an effective tool to count cycles. As illustrated in Fig. 2, a line segment is constructed in the state space from N_e to one edge of the triangle. According to dynamics models and the evolutionary stable state concept in EGT [9, 10, 25], we postulate that, if cycles exist, they should surround the center — the Nash equilibrium denoted as N_e at $(\frac{1}{3},\frac{1}{3})$. As mentioned above, each group would have a 300 rounds trajectory including 299 transits in state space. Some of the transits cross the Poincare section (S) from left to right (counter-clockwise) and some from right to left (clockwise).

We denote every transit-i as C_i, assign the transit-i crossing S as $C_i = 1$ (-1) when the transit is counter-clockwise (clockwise) and assign the transit-i not crossing S as $C_i = 0$. Then, we defined the number of the counted cycles in time interval $[t_0, t_1]$ as

$$C^e_{t_0,t_1} = \sum_i C_i, \tag{1}$$

If the evolutionary trajectory is full of random transits as expected by CGT, in long-run, C^e should be zero, otherwise, if $C^e > 0$ significantly, there should exist distance from equilibrium. Mathematically, C^e is of time odd $C^e_{t_1,t_0} = -C^e_{t_0,t_1}$, and will be discussed below.

Frequency of cycles —— To compare the cycles in different length of game, we adopt *frequency* as an index. Frequency denoted as f_e means the cycles in per experimental round and is calculated as

$$f_e = \frac{\sum_i C_i}{t_1 - t_0}. \tag{2}$$

This observable quantitatively captures the distance from equilibrium at one instantaneous snapshot in the experimental trajectories. The physical mean of the frequency is, on average, how many cycles can be obtained in each experimental round.

RESULTS

In all 12 300-rounds experiments, the strategies S, R and P have been used 6961, 7702 and 6937 times respectively. On average, the proportion of strategy S, R and P is 0.322, 0.357 and 0.321 respectively. This result is close to Nash equilibrium $(1/3, 1/3, 1/3)$.

There are total 12 trajectories obtained from the experiments. Indeed, the experimental trajectories are highly stochastic. This result is the same as the observations in other experiments [14, 15, 18, 20, 21]. Our main task is to test the cycles in the 12 experimental trajectories. The results are as follows.

Frequency of cycles —— Using Eq. 1, we calculate $C^e_{1,300}$. For each of the 12 experimental trajectories, the numerical results of $C^e_{1,300}$ are shown in the last column in Table I.

On average, over the 12 experimental trajectories, $\bar{C}^e_{1,300}$=8.50 ± 2.17. Accordingly, the observed frequency of cycles ($\bar{f}_e$) is 0.028 ± 0.007. That is to say, the empirical frequency of the cycles is 0.028 ±0.007 period per experimental round. And the 95% confidence interval is [0.012 0.044].

Existence of cycles—— To estimate the existence of the cycles, we use $C^e_{1,300}$ from the 12 group samples. Statistical result shows that H_1: $C^e_{1,300}$=0 can be rejected

TABLE I: C^e in the 12 trajectories

Group	$C^e_{1,50}$	$C^e_{1,100}$	$C^e_{1,150}$	$C^e_{1,200}$	$C^e_{1,250}$	$C^e_{1,300}$
1	1	4	6	6	11	11
2	-1	2	5	7	7	9
3	1	-1	5	8	12	17
4	1	2	6	8	11	11
5	-2	-2	-5	-4	-1	-3
6	3	2	2	7	9	12
7	3	4	12	17	20	23
8	2	5	7	10	11	12
9	-3	-5	-4	-3	0	0
10	0	1	5	3	5	5
11	-1	-1	-2	-1	-1	0
12	0	1	2	1	4	5
Mean	0.33	1.00	3.25	4.92	7.33	8.50
S.E.	0.54	0.83	1.42	1.73	1.82	2.17
p	0.55	0.25	0.04	0.02	0.00	0.00

The last three rows are the statistical results of the 12 experimental groups. The last row is p-value of the *t-test* by setting the null hypothesis $C^e_{1,t}$ of the 12 samples above.



FIG. 3: Counted cycles $C^e_{1,t}$ of the 12 sessions with t = 5, 10, 15, ..., 300. For example, ending at +23 at t = 300, the blue curve (the 7-*th* group in the 12 groups) reads 23 counter-clockwise cycles.

($p = 0.002 < 0.05$, *t-test*, two-tailed, $d.f.$=11). So, we can say that cycles do exist in the standard RPS game in discrete time experimental setting.

Direction of cycles —— Statistical result also shows $C^e_{1,300} > 0$ (p=0.001), that means the direction of cycles are of counter-clockwise. This result meets the direction expected by variety evolution dynamics [9, 10, 25]. Setting $t = 5, 10, 15, ..., 300$, the $C^e_{1,t}$ of each group are exhibited in Fig. 3. Nine out of twelve groups the counted cycles are positive, two groups are zero and one is negative; Statistically, the C^e is positive ($p < 0.01, ttest$ with 9 sample of +1, 2 samples of 0 and 1 sample of -1) and the direction is of counter-clock.

Persistence of cycles—— To see the persistence of cycles, we set the null hypothesis as H_2: $C^e_{1,150} = C^e_{151,300}$. If H_2 can be rejected in the data, cycles do persist, which mean EGT is supported and CGT is declined and vice versa. We use the p-values from *t-test* of the null hypothesis (the dissipation of cycles) H_2: $C^e_{1,150} = C^e_{151,300}$. Statistical result is $p > 0.10$. So, in our 12 300-rounds trajectories, the dissipation of cycles in the data can be rejected. In another word, the persistence of the cycles cannot be rejected by the data.

Accordingly, this result is our answer to the question 'Do cycles dissipate when subjects must choose simultaneously' raised by the authors [17].

Length of the experimental round to confirm cycles —— One result need to be emphasized. We have test, at what value of t, the null hypothesis H_1: $C^e_{1,t}$=0 can be rejected. According to the last row in Table I, we find that, the p-values are below 0.05 only when $t \geq 150$. This means that, using the 12 trajectories, to confirm the existence of cycles, the length should be *at least* 150 rounds. This point relates to the marginal value to test out cycles and will be discussed below.

MODEL EVALUATING

Till now, using frequency of cycles to index the distance from equilibrium, we have reported the empirical results from the experiments data. As evaluating parameters of dynamics models is a task in EGT investigation (e.g., [27]), in this section, we apply the frequency of cycles, as a novel way, to evaluate parameters in EGT.

For simple and referring to the experimental literatures to evaluate EGT [17–19], we use the continuous time and infinite population size approximation. Here, the demonstration is to evaluate the parameter in logit dynamics model in EGT. This dynamics works as follows: at every stage of the game players are selected uniformly at random according to a noisy best-response where the noise level is tuned by a parameter β [25, 28]. This parameter is a linking from the temperature in statistical physics to the degree of the rationality of human agents in strategy interaction [19, 29–32]. Explicitly, the logit dynamics can be presented as,

$$\dot{x_i} = \frac{\exp{(\beta U_i)}}{\sum_j \exp(\beta U_i)} - x_i, \tag{3}$$

in which U_i is the mean payoff of the agents using strategy i. The payoff-matrix of the game is in Fig. 1. At state (x, y, z), the mean payoff U_i of the three strategy can be presented as

$$\begin{bmatrix} U_x \\ U_y \\ U_z \end{bmatrix} = \begin{bmatrix} 2 - 2\,y - x \\ 2\,x + y \\ y - x + 1 \end{bmatrix} \tag{4}$$

It is not surprise that, the mean payoff ($\bar{U} = xU_x + yU_y + zU_z$) of this population always 1; Because the standard RPS game is a constant sum game and sum of each pair players are 1.

According to Eq. 4, the logit dynamics equation (Eq. 3) can be specified to

$$\begin{bmatrix} \dot{x} \\ \dot{y} \end{bmatrix} = \begin{bmatrix} \mathrm{e}^{\beta\,(2-x-2\,y)}X - x \\ \mathrm{e}^{\beta\,(2\,x+y)}X - y \end{bmatrix} \tag{5}$$

in which $X{=}\left[\mathrm{e}^{\beta\,(2\,x+y)} + \mathrm{e}^{\beta\,(y-x+1)} + \mathrm{e}^{\beta\,(2-x-2\,y)}\right]^{-1}$. As the density strategy of Scissors (z) always determined when the density strategy of Rock (x) and Paper (y) are given ($z = 1-x-y$), $\dot{z}$ is determined by $\dot{x}$ and $\dot{y}$ in Eq. 5 and can be ignored.

At a equilibrium state, the velocity of the strategy change should be 0. Then, setting the right-hand-side of Eq. 5 as 0, solving the equations, the equilibrium state can be obtained. That is to say, for a given β, we can obtain the mixed strategy equilibrium. No surprising, for *any* given β, (1/3, 1/3, 1/3) is always the solution and is the equilibrium state.

Surround the equilibrium state, the Jacobian is the function of β and can be expressed explicitly as

$$J = \begin{bmatrix} -\frac{1}{3}\beta - 1 & -\frac{2}{3}\,\beta \\ \frac{2}{3}\,\beta & \frac{1}{3}\beta - 1 \end{bmatrix} \tag{6}$$

From which, the eigenvalue ϵ of this Jacobian is

$$\epsilon = \ -1 \pm \tfrac{\sqrt{3}}{3}\,\beta\mathrm{i} \tag{7}$$

As the frequency of cycles is determined by the image part of the eigenvalue divided by 2π, so theoretical frequency of cycle $f_t(\beta) = \frac{\sqrt{3}}{6\pi}\,\beta$.

Using the empirical frequency (f_e), we have the best fit ($f_t = f_e$) for the model parameter β is,

$$\beta = 0.305 \pm 0.076 \tag{8}$$

and the 95% of C.I. of β is [0.131 0.479]. This parameter could index an experimental population (e.g., the degree of rationality [30]). The fixed parameter β in Eq. 8 could links to fundamental models in game theory [21, 30] and free energy like observations [1]. In this section, we have demonstrated that experiment test of non-equilibrium game behavior is possible.

DISCUSSION

In a standard RPS experiment with discrete time setting as [13–15, 18–21], we found the cycles exist and persist. Firstly, we test out the frequency of cycles in discrete time RPS game. We have also demonstrated how to fix the model parameter using the empirical frequency.

Comparing with existed experiments on EGT testing, we suggest that, our finding base on the two advantage points: (1) We have considered the time reversal asymmetry in measurement. (2) We have considered the sampling size effect in our experiment.

1. **Time reversal symmetry** —— As the central measurement of this report, C^e in Eq.(1) is of time odd ($C^e_{t_1,t_2}$=-$C^e_{t_2,t_1}$). In this way, the cycles beyond detailed balance can be counted. One can notice that, for the cycles counting in a stochastic trajectory, this measurement has its root in non-equilibrium statistical physics (e.g., see the observable $N_+ - N_-$ in [24]).

   Referring to existed experiment literatures, as emphasis by reference [17, 33], cycle in evolutionary trajectory has being sought for long time. Till now, in discrete time game, the existence of cycles is only supported qualitatively [10, 14, 18] but not quantitatively.

   Recently, two time odd observations, velocity vector [26] and angular momentum [34], are developed to indexed the experimental evolutionary trajectory. Both of the observations can test out EGT patterns in significant. No surprising, these time odd observable, like entropy production [24, 35–37], can capture the excess transits beyond detailed balance and can index the distance from equilibrium in games.

2. **Sample size effect** —— To test out the deterministic behaviors, in higher stochastic system, longer experimental rounds and groups numbers is needed. It is sample size effect. This picture has been demonstrated in the experiment in non-equilibrium statistical physics [2]. For example, the Fig. 1 in ref. [2], using 540 experimental trajectories, the authors compare two time interval conditions (10^{-2} sec and 2 sec with sampling $\triangle t = 10^{-3}$ sec). For 10^{-2} sec, the probability of a trajectory and its time reversal trajectory seems nearly equal. On the contrast, the asymmetry in 2 sec samples can be test out significantly. In experimental game trajectories, the condition is similar.

   As large sample size are appropriate, we set the experimental round up to 300 rounds. According to the statistical result in our data, $C^e_{1,t} > 0$ could be significant only when $t \geq 150$ (see the last paragraph in the Result section). That is to say, even using 12 experimental trajectories, if the experimental rounds are less than the margin, we cannot confirm the existence of cycles even though time odd measurement has been used as Eq. 1. That is also the reason why we set our experiments up to 300 rounds.

   Referring to related discrete time experiments on EGT, the number of rounds and the groups usually are less. For example, in the experiments [18], the experimental rounds is 100 and the tested groups is 4 for a given parameter; As a result [17], cycles cannot be test out. Usually, the experimental trajectories are less than 12 and the experimental rounds are less than 100 in a trajectory, from these experiments, to confirm the existence of cycles should be uneasy.

Accordingly, we suggests that, for further investigating the cycles in experiments, longer experiments rounds and larger experimental trajectories are expected. Only when the sampling size is large enough, excess observations beyond detailed balance can be statistically significant.

It can be noticed that both of the two points are rooting in non-equilibrium statistical physics and for measuring the distance from equilibrium.

Linking experimental economics game to principles in statistical physics is developing, e.g., maximum entropy principle governs Nash equilibrium [38] and governs social transits [39]; In lab game experiments [36, 37], fluctuation relations (Fluctuation Theorem) [24, 40] is supported. Evidences from experimental economics promote that there could exist wider common background of social science and physical science [41].

SUMMARY

In standard RPS game, surround Nash equilibrium, EGT predicts cycles; Alternatively, CGT predict no cycle exists and the evolutionary trajectories stochastic. Results from our traditional setting (discrete time and random matching) experiments show that, the evolutionary trajectories are high stochastic indeed, however, surround Nash equilibrium, cycles exist and persist. Interestingly, we test out the frequency of cycles. As demonstrated, the empirical frequency could be linked the evolutionary dynamics model. Our measurement has the roots in non-equilibrium statistical physics.

We wish, basing on game theory and non-equilibrium statistical physics, the co-playing of lab experiments and dynamics models could lead us to see something *what we have not seen yet.*

Acknowledgements —— We thank Ken Binmore for helpful discussion. This work is supported by the grant from State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences and the grant from Experimental Social Science Laboratory of Zhejiang University. We also thank Zunfeng Wang and Shuang Wang for assistance.

---

[1] D. Sivak and G. Crooks, Physical Review Letters **108**, 150601 (2012).

[2] G. Wang, E. Sevick, E. Mittag, D. Searles, and D. Evans, Physical Review Letters **89**, 50601 (2002).

[3] N. Kumar, S. Ramaswamy, and A. Sood, Physics Review Letters **106**, 118001 (2011).

[4] E. Frey, Physica A: Statistical Mechanics and its Applications **389**, 4265 (2010).

[5] B. Sinervo and C. Lively, Nature **380**, 240 (1996).

[6] B. Kirkup and M. Riley, Nature **428**, 412 (2004).

[7] D. Friedman, Journal of Evolutionary Economics **8**, 423 (1998).

[8] J. Von Neumann and O. Morgenstern, *Theory of Games and Economic Behavior* (Princeton university press, 1944).

[9] J. Smith, *Evolution and the Theory of Games* (Cambridge university press, 1982).

[10] M. Hoffman, S. Suetens, M. Nowak, and U. Gneezy, Science, submit (2012).

[11] G. N. Lewis, Proceedings of the National Academy of Sciences **11**, 179 (1925).

[12] A. Falk and J. Heckman, Science **326**, 535 (2009).

[13] L. Samuelson, The Journal of Economic Perspectives **16**, 47 (2002).

[14] K. Binmore, J. Swierzbinski, and C. Proulx, The Economic Journal **111**, 445 (2001).

[15] Y. Cheung and D. Friedman, Journal of economic behavior and organization **35**, 263 (1998).

[16] D. Semmann, H.-J. Krambeck, and M. Milinski, Nature **425**, 390 (2003), 10.1038/nature01986.

[17] T. Cason, D. Friedman, and E. Hopkins, *Cycles and instability in a rock-paper-scissors population game: a continuous time experiment* (2012).

[18] D. Friedman, The Economic Journal **106**, 1 (1996).

[19] R. Battalio, L. Samuelson, and J. Van Huyck, Econometrica **69**, 749 (2001).

[20] J. Van Huyck, Handbook of Experimental Economics Results **1**, 520 (2008).

[21] R. Selten and T. Chmura, The American Economic Review **98**, 938 (2008).

[22] D. Vilone, A. Robledo, and A. Sánchez, Physical review letters **107**, 038101 (2011), pRL.

[23] J. Weibull, *Evolutionary game theory* (The MIT press, 1997).

[24] D. ben Avraham, S. Dorosz, and M. Pleimling, Physical Review E **84**, 011115 (2011), pRE.

[25] W. Sandholm, *Population games and evolutionary dynamics* (MIT press Cambridge, MA:, 2011).

[26] B. Xu and Z. Wang, *Evolutionary Dynamical Pattern of ”Coyness and Philandering”: Evidence from Experimental Economics*, vol. VIII (p1313-1326, NECSI Knowledge Press, ISBN 978-0-9656328-4-3., 2011).

[27] A. Traulsen, D. Semmann, R. D. Sommerfeld, H.-J. Krambeck, and M. Milinski, Proceedings of the National Academy of Sciences **107**, 2962 (2010).

[28] J. Hofbauer and K. Sigmund, Bulletin of the American Mathematical Society **40**, 479 (2003).

[29] L. Blume et al., Games and economic behavior **5**, 387 (1993).

[30] C. Watkins and P. Dayan, Machine learning **8**, 279 (1992).

[31] R. McKelvey and T. Palfrey, Games and economic behavior **10**, 6 (1995).

[32] A. Kianercy and A. Galstyan, Physical Review E **85**, 041145 (2012).

[33] M. Benaīm, J. Hofbauer, and E. Hopkins, Journal of Economic Theory **144**, 1694 (2009), ISSN 0022-0531.

[34] Z. Wang and B. Xu, arXiv preprint arXiv:1203.2591

(2012).
[35] B. Andrae, J. Cremer, T. Reichenbach, and E. Frey, Physical review letters **104**, 218102 (2010).
[36] B. Xu and Z. Wang, Available at SSRN 1878063 (2011).
[37] B. Xu and Z. Wang, Available at SSRN 1988985 (2012).
[38] Physics Letters A **376**, 1318 (2012).
[39] B. Xu and Z. Wang, Results in Physics **2**, 127 (2012).
[40] D. Evans, E. Cohen, and G. Morriss, Physical Review Letters **71**, 2401 (1993).
[41] D. H. Wolpert, M. Harre, E. Olbrich, N. Bertschinger, and J. Jost, Physical Review E **85**, 036102 (2012).